

Ministry of Education and Science of the Russian Federation
Saint Petersburg National Research University of Information
Technologies, Mechanics, and Optics

NANOSYSTEMS:

PHYSICS, CHEMISTRY, MATHEMATICS

2013, volume 4(4)

Special Issue

**INTERNATIONAL CONFERENCE
"MATHEMATICAL CHALLENGE OF QUANTUM
TRANSPORT IN NANOSYSTEMS - 2013.
PIERRE DUCLOS WORKSHOP"**

Наносистемы: физика, химия, математика

2013, том 4, № 4



NANOSYSTEMS:

PHYSICS, CHEMISTRY, MATHEMATICS

ADVISORY BOARD MEMBERS

Chairman: V.N. Vasiliev (*St. Petersburg, Russia*),
V.M. Ievlev (*Voronezh*), P.S. Kop'ev (*St. Petersburg, Russia*),
V.N. Parmon (*Novosibirsk*), A.I. Rusanov (*St. Petersburg, Russia*).

EDITORIAL BOARD

Editor-in-Chief: N.F. Morozov (*St. Petersburg, Russia*)
Deputy Editor-in-Chief: I.Yu. Popov (*St. Petersburg, Russia*)

Section Co-Editors:

Physics – V.M. Uzdin (*St. Petersburg, Russia*),
Chemistry, material science – V.V. Gusarov (*St. Petersburg, Russia*),
Mechanics – A.K. Belyaev (*St. Petersburg, Russia*),
Mathematics – I.Yu. Popov (*St. Petersburg, Russia*).

Editorial Board Members:

V.M. Adamyan (*Odessa, Ukraine*); O.V. Al'myasheva (*St. Petersburg, Russia*); S. Bechta (*Stockholm, Sweden*); V.G. Bepalov (*St. Petersburg, Russia*); A. Chatterjee (*Hyderabad, India*); S.A. Chivilikhin (*St. Petersburg, Russia*); A.V. Chizhov (*Dubna, Russia*); P.P. Fedorov (*Moscow, Russia*); E.A. Gudilin (*Moscow, Russia*); D.A. Indeitsev (*St. Petersburg, Russia*); A.L. Ivanovskii (*Ekaterinburg, Russia*); H. Jónsson (*Reykjavik, Iceland*); A.A. Kiselev (*Madison, USA*); Yu.S. Kivshar (*Canberra, Australia*); S.A. Kozlov (*St. Petersburg, Russia*); Yu.B. Kudasov (*Sarov, Russia*); S.E. Kul'kova (*Tomsk, Russia*); P.A. Kurasov (*Stockholm, Sweden*); A.V. Lukashin (*Moscow, Russia*); V.A. Margulis (*Saransk, Russia*); I.V. Melikhov (*Moscow, Russia*); G.P. Miroshnichenko (*St. Petersburg, Russia*); H. Neidhardt (*Berlin, Germany*); K. Pankrashkin (*Orsay, France*); B.S. Pavlov (*Auckland, New Zealand*); A.V. Ragulya (*Kiev, Ukraine*); V. Rajendran (*Tamil Nadu, India*); V.P. Romanov (*St. Petersburg, Russia*); V.Ya. Rudyak (*Novosibirsk, Russia*); V.M. Talanov (*Novocherkassk, Russia*); A.Ya. Vul' (*St. Petersburg, Russia*).

Editors:

I.V. Blinova; A.I. Popov; M.A. Skryabin; A.I. Trifanov; E.S. Trifanova (*St. Petersburg, Russia*),
R. Simoneaux (*Philadelphia, Pennsylvania, USA*).

Address: University ITMO, Kronverkskiy pr., 49, St. Petersburg 197101, Russia.

Phone: +7(812)232-67-65, **Journal site:** <http://nanojournal.ifmo.ru/>,

E-mail: popov1955@gmail.com

AIM AND SCOPE

The scope of the journal includes all areas of nano-sciences. Papers devoted to basic problems of physics, chemistry, material science and mathematics inspired by nanosystems investigations are welcomed. Both theoretical and experimental works concerning the properties and behavior of nanosystems, problems of its creation and application, mathematical methods of nanosystem studies are considered.

The journal publishes scientific reviews (up to 30 journal pages), research papers (up to 15 pages) and letters (up to 5 pages). All manuscripts are peer-reviewed. Authors are informed about the referee opinion and the Editorial decision.

Content

From the Editorial Board 445

**INTERNATIONAL CONFERENCE
"MATHEMATICAL CHALLENGE OF QUANTUM TRANSPORT
IN NANOSYSTEMS - 2013. PIERRE DUCLOS WORKSHOP"**

INVITED SPEAKERS

Ibrahim Baydoun, Valentin A. Zagrebnov
Diffusion and laplacian transport for absorbing domains 446

Mikkel H. Brynildsen , Horia D. Cornean
On spectral gaps in graphene in a weak constant magnetic field 467

Konstantin Pankrashkin
**On the asymptotics of the principal eigenvalue for a Robin problem
with a large parameter in planar domain** 474

M.I. Belishev
C*-algebras in reconstruction of manifolds 484

CONTRIBUTED TALKS

J. Smotlacha, R. Pincak
**Application of Green's function approach to electronic structure
of carbon nanocylinders** 490

E. Karpov, Schafer, O.V. Pilyavets, R. Garcia-Patron, N. J. Cerf
Gaussian classical capacity of gaussian quantum channels 496

D.R. Gulevich, P.N. Dmitriev, V.P. Koshelets, F.V. Kusmartsev
**Experimental and Theoretical Characterisation of Josephson
Self-coupling in Sub-Terahertz Flux-flow Oscillator** 507

I. S. Lobanov, A. I. Trifanov, E. S. Trifanova
**Genetic algorithm for constructing graphene nanoribbon with given
electronic transport properties** 512

I.A. Gladskikh, N.B. Leonov, S.G. Przhibel'skii, T.A. Vartanyan
Hysteresis of conductivity in the granular silver films 524

Yu.Kh. Eshkabilov, R.R. Kucharov
Efimov's effect for partial integral operators of Fredholm type 529

Natalie E. Firsova
The conductivity low energy asymptotics for monolayer graphene 538

YOUNG RESEARCHES SIMPOSIUM

A.S. Anikevich
Negative eigenvalues of the Y-type chain of weakly coupled ball resonators 545

A.E. Ivanova, V.I. Egorov, S.A. Chivilikhin, A.V. Gleim
Investigation of quantum random number generation based on space-time division of photons 550

N.N. Konobeeva, M.B. Belonenko
Tunneling current in carbon nanotubes with deep impurities 555

I.F. Melikhov, I.Yu. Popov
Hartree-Fock approximation for the problem of particle storage in deformed nanolayer 559

A.V. Pak, M.B. Belonenko, P.N. Dyachkov
Electrophysical properties of CNT. The augmented cylindrical wave method 564

K.V. Pravdin, I.Yu. Popov
Model of the interaction of point source electromagnetic fields with metamaterials 570

Information for authors 577

FROM THE EDITORIAL BOARD

In this issue we publish the Proceedings of the International Conference **Mathematical Challenge of Quantum Transport in Nanosystems**. “**Pierre Duclos Workshop**” organized by Saint Petersburg National Research University of Information Technologies, Mechanics, and Optics in March 2013. This workshop was the eighth of a series which has the aim to bring together specialists in nanosystem modeling, mathematicians and condensed matter physicists

PROGRAM COMMITTEE

Vladimir N. Vasilyev (University ITMO, Russia) – Chairman
Igor Yu. Popov (University ITMO, Russia) – Vice-Chairman
Sergey N. Naboko (St. Petersburg State University, Russia)
Boris S. Pavlov (Massey University, New Zealand)
George P. Miroshnichenko (University ITMO, Russia)

LOCAL ORGANIZING COMMITTEE

Leonid M. Studenikin – Chairman
Igor Yu. Popov – Vice-Chairman
Igor S. Lobanov
Ekaterina S. Trifanova
Irina V. Blinova
Alexander I. Trifanov
Maxim A. Skryabin
Anton I. Popov

INVITED SPEAKERS

Pavel Exner (Prague, Czech Republic)
Horia Cornean (Aalborg, Denmark)
Hagen Neidhardt (Berlin, Germany)
Valentin Zagrebnov (Marseille, France)
Konstantin Pankrashkin (Paris, France)
Mikhail Belishev (St. Petersburg, Russia)
Jaroslav Dittrich (Rez, Czech Republic)
Paul Tackley (Zurich, Switzerland)

THE MAIN TOPICS OF THE CONFERENCE:

Spectral theory
Scattering
Quantum transport
Quantum communications and computations

ELECTROPHYSICAL PROPERTIES OF CNT. THE AUGMENTED CYLINDRICAL WAVE METHOD

A. V. Pak¹, M. B. Belonenko^{2,3}, P. N. Dyachkov⁴

¹ Volgograd State University, 400062, Volgograd, Russia

² Entropique Inc., London, N6J 3S2, ON, Canada

³ Volgograd Institute of Business, 400048, Volgograd, Russia

⁴ Institute of General and Inorganic Chemistry named after N.C. Kurnakov RAS,
Moscow, Russia

pak.anastasia@gmail.com

PACS 71.10.Fd, 72.80.Vp

Current-voltage characteristics of tunnel contact between semiconducting (and conducting) carbon nanotubes (CNT) of various diameters and system of periodically located quantum dots (and also in contact to metal) was obtained using density of states (DOS) investigation. DOS has been calculated by means of the method of attached cylindrical waves. At certain parameters for quantum dots, the current-voltage characteristics observed testify to the presence of negative differential conductivity.

Keywords: carbon nanotubes, density of the states, current-voltage characteristics.

1. Introduction

In the last few years, the interest of many researchers has been focused on the properties of carbon nanotubes [1]. It is known from classical works that the “Arm-chair” type of nanotube, in Huckel’s approximation, are conductors at any temperature [2]. However, using the linear augmented cylindrical wave (LACW) method allows a gap in the band structure of this type of nanotube. This makes it possible to use them in different applications. The simplest method to experimentally verify the change in the state density is to study the tunneling current [3,4], for example, of the contact with quantum dots. In these papers, the density of states were calculated by the method of Green’s functions [5], however, not every system has a density of states for electrons can be determined analytically. For such a system, the more versatile method is based on ab initio calculations .

Theoretically, investigation was done by tight-binding calculations only, but it is known from the band structure theory of solids that the linear combination of atomic orbitals (LCAO) basis is adequate to achieve good results for the valence band, but not for the conduction band. The reason for this is that this basis does not include the delocalized conducting plane-wave type functions. As to the π electron band structure models, they are adequate for calculating the energy curves located in a Fermi level region only. Therefore, we have calculated the complete band structures of metallic armchair (n, n) CNTs with $4 < n < 100$ and zigzag semiconducting $(0, n)$ CNTs with $10 < n < 49$ indivisible by 3 in terms of a LACW method. The LACW method is just a reformulation of the linear augmented plane wave (LAPW) theory for cylindrical multiatomic systems. Its basis functions have both localized and delocalized components. Finally, the main argument for using cylindrical waves is to account for the cylindrical geometry of the nanotubes in an explicit form that offers obvious advantages [6].

2. Basic equations

2.1. Cylindrical muffin-tin potential

Similar to the LAPW technique used in the band structure theory of solids, we apply the muffin-tin (MT) and local density functional approximations for electronic potentials of tubules. However, the potential of a nanotube differs drastically from that of bulk material. Indeed, infinite motion of an electron is possible in any direction in a crystal, but it is obviously limited in the case of nanotubes by their size and cylindrical shape. Therefore, in terms of the LACW method, the atoms of nanotube are considered to be enclosed between two impenetrable cylinder-shaped potential barriers Ω_a and Ω_b , because there are two vacuum regions Ω_v on the outside and on the inside of the tubule. The radii a and b of these barriers are chosen so that the region confined by barriers accommodates a significant portion of the electron density of the tubule. Based on our previous calculations of the CNTs, we take $a = R_{NT} + 2.3$ a.u., $b = R_{NT} + 2.3$ a.u., where, R_{NT} is radius of the tubule. Finally, the electronic potential is spherically symmetrical in the regions of MT-spheres of atoms α and constant in the interspherical region. Inside these spheres, we calculate the electron potential by means of the local density approximation with Slater exchange. As usually, the radii of the MT spheres were chosen so that the atomic spheres touch, but do not overlap. Outside the MT spheres up to the impenetrable potential barriers Ω_a and Ω_b , the potential is approximated by a constant value taken to be the energy zero point [6,7].

2.2. Solution of the Schrödinger equation

The basis functions called LACWs are solutions of the Schrödinger equation for the interspherical and MT regions of tubule sewn together so that the resulting LACWs are continuous and differentiable anywhere in the system. In the interspherical region, the LACWs are the solutions of the Schrödinger equation for free electron movement in the infinitely long potential well between two impenetrable cylindrical potential barriers. Here, the solutions are the superpositions of the cylindrical Bessel functions of the first and second kinds. In any MT sphere, the solutions are expanded in terms of the spherical harmonic functions and the eigenfunctions of the radial Schrödinger equation. To obtain the basis wave functions, the solutions of Schrödinger equation in these two regions are matched on the surfaces of the MT spheres so that both the basis functions and their normal derivatives are continuous across these surfaces. The eigenfunctions of electrons in the tubule are the linear superpositions of these basis LACWs. The coefficients of the expansions of any eigenfunction in terms of these basis functions and the electron dispersion curves are defined by the variational method.

In this work, we apply the symmetry-adapted version of the LACW method developed previously. In this version, one takes into account that every single-walled CNT can be generated by first mapping only two nearest-neighbor C atoms onto a surface of a cylinder and then using the rotational and helical symmetry operators to determine the remainder of the tubule. With account of these symmetries, the cells contain only two carbon atoms, and the *ab initio* LACW theory becomes applicable to any tubule independent of the number of atoms in a translational unit cell. In the LACW studies, we use only this structural information as input parameters.

The dispersive relation for quantum wells is similar to the band structure of superlattice [8]:

$$E_{QD} = \varepsilon_0 - \Delta \cos(p),$$

where ε_0 is electron energy in quantum well, Δ is tunnel integral, defining by overlap of electron wave functions of nearest wells.




FIG. 1. Density of states for CNT arm-chair type: a) Huckel's approximation; b) LACW method

The density of states for arm-chair CNTs, calculated under the π -electron approximation [2], was shown in Fig. 1a. The LACW method allows in the calculation consideration of more deep-seated σ -orbital electrons at very high temperatures (~ 25 – 27 eV) to get a gap in the energy spectrum (see Fig. 1b). This fact allowed the calculation of the electrical properties of conductive nanotubes in contact with the quantum dots.




FIG. 2. Current-voltage characteristics for CNT type "zig-zag" of various diameter ((10, 0) – (50, 0)) in contact with quantum dots system

3. Tunnel characteristics

In order to study tunneling effects, we will set Hamiltonian of our model in a form:

$$H = \sum_p \varepsilon_p^A a_p^+ a_p + \sum_q \varepsilon_q^B b_q^+ b_q + \sum_{pq} T_{pq} (a_p^+ b_q + b_q^+ a_q), \quad (1)$$

where a_p^+ , a_p - production (annihilation) operators of electrons with a momentum p in carbon nanotubes; ε_p^A - electron energy spectrum of nanotubes (1); T_{pq} - matrix element of the operator of tunneling between conditions p and q ; b_q^+ , b_q - creation (annihilation) operators of electrons with an momentum q in matter, resulted in contact with carbon nanotubes; ε_p^B - electron energy spectrum of quantum dots. Note, that in (1) p and q are multi-indices. Also, if we consider the external electric field \vec{E} which we will investigate using the gauge: $\vec{E} = -\frac{1}{c} \frac{\partial \vec{A}}{\partial t}$, it is possible to take it into account by means of replacement the corresponding momentum components: $p \rightarrow p - eA/c$.

Defining a tunnel current as [5]:

$$J = ie \sum_{pq} (a_p^+ b_q - b_q^+ a_p), \quad (2)$$

and carrying out calibration transformation [5]:

$$a_p \rightarrow S^{-1} a_p S, \\ S = \exp \left(ieVt \sum_p a_p^+ a_p \right),$$

where V applied, for definiteness, to CNT voltage, e - electron charge, it is possible to reduce formally a problem about calculation characteristic current-voltage to calculation of the response of the operator: $J_t = ie \sum_{pq} (a_p^+ b_q e^{ieVt} - b_q^+ a_q e^{-ieVt})$ on external action [5]:

$$H_t = \sum_{pq} T_{pq} (a_p^+ b_q e^{ieVt} + b_q^+ a_q e^{-ieVt}).$$

Within the framework of Kubo theory, the answer is set by the formula:

$$J = 4\pi e |T|^2 \int_{-\infty}^{\infty} d\varepsilon \nu_A(\varepsilon + eV) \nu_B(\varepsilon) (n_f(\varepsilon) - n_f(\varepsilon + eV)), \quad (3) \\ \nu_A(\varepsilon) = \sum_p \delta(\varepsilon - \varepsilon_p^A), \quad \nu_B(\varepsilon) = \sum_q \delta(\varepsilon - \varepsilon_q^B),$$

where $\delta(x)$ - Dirac delta-function, $\nu_{A(B)}(\varepsilon)$ - tunnel density of states; $n_f(\varepsilon)$ - equilibrium number of fermions with energy ε . We used the approach of "rough" contact: $T_{pq} = T$ (a nanotube is perpendicular to the quantum dot surface). After integrals calculation entering in (3) it is easy to obtain current-voltage characteristic of the contact presented in Fig. 2–3.

The existence of a descending section of the current-voltage characteristic of the tunneling conductance between the nanotubes and the quantum dots system can be clearly seen in Fig. 2. This indicates the presence of negative differential conduction. The descending section is observed at the same parameters of quantum dots: $\varepsilon_0 = 22$ eV - energy of electrons a quantum




FIG. 3. Current-voltage characteristics for CNT type “arm-chair” of various diameter ((4,4) – (9,9)) in contact with quantum dots system

hole, and $\Delta = 3$ eV - the tunnel integral defined by overlapping of wave electronic functions of the next holes. The size of the resistance obtained in such a system was $R \approx -24$ kOhm.

Similarly, the current-voltage characteristics for semiconducting CNT of various diameters ((10,0)–(50,0)) in contact with systems of quantum dots are represented in Fig. 2. For semiconducting CNT in contact with a quantum dots system, the descending section in the current-voltage, characteristic of tunnel conductivity, was also observed. The resistance obtained for such a system was $R \approx -18$ kOhm (should be -18 kOhm) for the same parameters of the quantum dots system.

It should be noted that the descending section in CVC for system CNT-QD is absent in the case of pi-approximation for CNT DOS.

The obtained dependencies can have the important practical applications for studying noncontact and the design of tunneling diodes based on carbon nanotubes.

4. The conclusion

In summary we will formulate the basic conclusions from the made work.

- (1) Current-voltage characteristics of contacts CNT - a superlattice of quantum dots are obtained.
- (2) The descending section with negative differential conduction is observed for certain characteristics of quantum dots. This effect can be used to create various frequency amplifiers.

References

- [1] E.G. Rakov. Carbon nanotubes in new materials. *Russian Chemical*, **82** (1), P. 27–47 (2013).
- [2] P. Harris. Carbon nanotubes and related structures. *New materials of XXI centuries*. Tehnosfera, Moscow: 2003. 336 p.
- [3] M.B. Belonenko, N.G. Lebedev, A.V. Pak. Tunneling current of the contact impurity Graphene nanoribbon - quantum dots. *Russian Physics Journal*, **55** (6), P. 644–648 (2012).

- [4] M.B. Belonenko, A.V. Pak, N.G. Lebedev. Tunnel current between impurity graphene nanoribbons. *Semiconductors*, **47** (5), P. 649–651 (2013).
- [5] L.S. Levitov, A.V. Shitov. *Green's functions. Problems with solutions* [In Russian]. Fizmatlit, oscow: 2003, 392 p.
- [6] P.N. D'yachkov, H. Hermann, D.V. Kirin. Electronic structure and interband transitions of metallic carbon nanotubes. *Applied Physics Letters*, **81**, P. 5228–5231 (2002).
- [7] P.N. D'yachkov, O.M. Kepp. Science and Application of Nanotubes, D. Tomanek Ed. NY, Plenum, P. 77–82 (2000).
- [8] A.P. Silin. Semiconductive superlattice [In Russian]. *Physics-Uspeski*, **147** (3), P. 485–516 (1985).



NANOSYSTEMS:

PHYSICS, CHEMISTRY, MATHEMATICS

INFORMATION FOR AUTHORS

The journal publishes research articles and reviews, and also short scientific papers (letters) which are unpublished and have not been accepted for publication in other magazines. Articles should be submitted in English. All articles are reviewed, then if necessary come back to the author to completion.

The journal is indexed in Chemical Abstract Service of the American Chemical Society and in Russian Scientific Citation Index.

Author should submit the following materials:

1. Article file in English, containing article title, the initials and the surname of the authors, Institute (University), postal address, the electronic address, the summary, keywords, MSC or PACS index, article text, the list of references.
2. Files with illustrations, files with tables.
3. The covering letter in English containing the article information (article name, MSC or PACS index, keywords, the summary, the literature) and about all authors (the surname, names, the full name of places of work, the mailing address with the postal code, contact phone number with a city code, the electronic address).
4. The expert judgement on possibility of publication of the article in open press (for authors from Russia).

Authors can submit a paper and the corresponding files to the following addresses: nanojournal.ifmo@gmail.com, popov1955@gmail.com.

Text requirements

Articles should be prepared with using of text editors MS Word or LaTeX (preferable). It is necessary to submit a pdf copy. In the name of files the English alphabet is used. The recommended size of short communications (letters) is 4-6 pages, research articles – 6-15 pages, reviews – 30 pages.

Recommendations for text in MS Word:

Formulas should be written using Math Type. Figures and tables with captions should be inserted in the text. Additionally, authors present separate files for all figures and Word files of tables.

Recommendations for text in LaTeX:

Please, use standard LaTeX without macros and additional style files. The list of references should be included in the main LaTeX file. Single LaTeX file of the paper with the corresponding pdf file should be submitted.

References in the article text are given in square brackets. The list of references should be prepared in accordance with the following samples:

- [1] N. Surname. *Book Title*. Nauka Publishing House, Saint Petersburg, 281 pp. (2000).
- [2] N. Surname, N. Surname. Paper title. *Journal Name*, **1** (5), P. 17-23 (2000).
- [3] N. Surname, N. Surname. Lecture title. In: Abstracts/Proceedings of the Conference, Place and Date, P. 17-23 (2000).
- [4] N. Surname, N. Surname. Paper title (2010). URL: <http://books.ifmo.ru/ntv>.
- [5] N. Surname, N. Surname. Patent Name. Patent No. 11111, Bul. No. 33, 5 pp. (2010).
- [6] N. Surname, N. Surname. Thesis Title. Thesis for full doctor degree in math. and physics, Saint Petersburg, 105 pp. (2000).

Requirements to illustrations

Illustrations should be submitted as separate black-and-white files. Formats of files – jpeg, eps, tiff.



NANOSYSTEMS:

PHYSICS, CHEMISTRY, MATHEMATICS

Журнал зарегистрирован

Федеральной службой по надзору в сфере связи, информационных технологий и массовых коммуникаций

(свидетельство ПИ № ФС 77 - 49048 от 22.03.2012 г.)

ISSN 2220-8054

Учредитель: федеральное государственное бюджетное образовательное учреждение высшего профессионального образования «Санкт-Петербургский национальный исследовательский университет информационных технологий, механики и оптики»

Издатель: федеральное государственное бюджетное образовательное учреждение высшего профессионального образования «Санкт-Петербургский национальный исследовательский университет информационных технологий, механики и оптики»

Отпечатано в типографии «Фалкон Принт»

Адрес: 197101, г. Санкт-Петербург, ул. Большая Пушкарская, 54

Подписка на журнал НФХМ

На 2013 год подписка осуществляется через ОАО Агентство «Роспечать»

Подписной индекс 57385 в каталоге «Издания органов научно-технической информации»